

Linear Programming: Duality

Daniel Kane

Department of Computer Science and Engineering
University of California, San Diego

Advanced Algorithms and Complexity
Data Structures and Algorithms

Learning Objectives

- Write the dual program of a linear program.
- Understand the duality theorem.

Example

Recall our first example:

Maximize $200M + 100W$ subject to:

- $W \geq 0$.
- $100 \geq M \geq 0$.
- $W \geq 2M$.
- $100,000 \geq 200(W - 2M) + 600M$.

Upper Bound

The best you could do was 60000, but we proved it by combining constraints

$$\begin{array}{rcl} 100 \cdot [& 001 \cdot M + 000 \cdot W & \leq 100] \\ +0.5 \cdot [& 200 \cdot M + 200 \cdot W & \leq 100,000] \\ \hline & 200 \cdot M + 100 \cdot W & \leq 60,000. \end{array}$$

General Technique

Try to prove bound by combining the constraints together.

Linear Program

Say you have the linear program where you want to minimize

$$v_1x_1 + v_2x_2 + \dots + v_nx_n$$

subject to constraints

$$a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n \geq b_1$$

...

$$a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n \geq b_m$$

Combine Constraints

If we have $c_j \geq 0$, we can combine constraints:

$$c_1 \cdot [a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n \geq b_1]$$

...

$$+ c_m \cdot [a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n \geq b_m]$$

$$w_1x_1 + w_2x_2 + \dots + w_nx_n \geq t,$$

$$w_i = \sum c_j a_{ji}, \quad t = \sum c_j b_j.$$

Bound

If $w_i = v_i$ for all i , have

$$v_1x_1 + v_2x_2 + \dots + v_nx_n \geq t.$$

Bound

If $w_i = v_i$ for all i , have

$$v_1x_1 + v_2x_2 + \dots + v_nx_n \geq t.$$

Want to find $c_j \geq 0$ so that $v_i = \sum_{j=1}^m c_j a_{ji}$ for all i , and $t = \sum_{j=1}^m c_j b_j$ is as large as possible.

Linear Program

Note that this is another **linear program**.

Find $c \in \mathbb{R}^m$ so that $\sum_{j=1}^m c_j b_j$ is as large as possible, subject to the linear inequalities $c_j \geq 0$, and equalities

$$v_i = \sum_{j=1}^m c_j a_{ji}.$$

Dual Program

Definition

Given the linear program (the primal):

Minimize $v \cdot x$

Subject to $Ax \geq b$

The **dual linear program** is the linear program:

Maximize $y \cdot b$

Subject to $y^T A = v$, and $y \geq 0$.

Bounds

It is not hard to show that a solution to the dual bounds the optimum for the primal.

Bounds

It is not hard to show that a solution to the dual bounds the optimum for the primal.

If $y \geq 0$ and $y^T A = v$,

then for any x with $Ax \geq b$,

$$x \cdot v = y^T Ax \geq y^T b = y \cdot b.$$

Bounds

It is not hard to show that a solution to the dual bounds the optimum for the primal.

If $y \geq 0$ and $y^T A = v$,
then for any x with $Ax \geq b$,

$$x \cdot v = y^T Ax \geq y^T b = y \cdot b.$$

The surprising thing is that these two linear programs always have the **same** solution.

Duality

Theorem

A linear program and its dual always have the same (numerical) answer.

Duality

Theorem

A linear program and its dual always have the same (numerical) answer.

This means that one can instead solve the dual problem. This is sometimes easier, and often provides insight into the solution.

Example: Flows

The size of our flow is

$$\sum_{\text{e out of a source}} f_e - \sum_{\text{e into a source}} f_e.$$

Example: Flows

The size of our flow is

$$\sum_{\text{e out of a source}} f_e - \sum_{\text{e into a source}} f_e.$$

By adding multiples of the conservation of flow equation, this is

$$\sum_v c_v \left(\sum_{\text{e out of } v} f_e - \sum_{\text{e into } v} f_e \right)$$

where $c_s = 1, c_t = 0$.

Example: Flows

This is

$$\sum_{e=(v,w)} (c_v - c_w) f_e.$$

Example: Flows

This is

$$\sum_{e=(v,w)} (c_v - c_w) f_e.$$

We can bound this using capacity constraints as

$$\sum_{e=(v,w)} C_e \max(c_v - c_w, 0).$$

Example: Flows

It is not hard to show that minimum attained when $c_v \in \{0, 1\}$.

Example: Flows

It is not hard to show that minimum attained when $c_v \in \{0, 1\}$.

Letting \mathcal{C} , be the set of vertices where $c_v = 1$, our bound is

$$\sum_{e=(v,w), v \in \mathcal{C}, w \notin \mathcal{C}} C_e = |\mathcal{C}|.$$

Example: Flows

It is not hard to show that minimum attained when $c_v \in \{0, 1\}$.

Letting \mathcal{C} , be the set of vertices where $c_v = 1$, our bound is

$$\sum_{e=(v,w), v \in \mathcal{C}, w \notin \mathcal{C}} C_e = |\mathcal{C}|.$$

The dual program just finds the minimum cut!

Example: Diet Problem

Recall the diet problem:

- Minimize the cost of the foods you need to buy subject to:
- Meets daily requirements for various nutrients
- Non-negative amount of each type of food

Example: Diet Problem

Recall the diet problem:

- Minimize the cost of the foods you need to buy subject to:
- Meets daily requirements for various nutrients
- Non-negative amount of each type of food

What is the dual program?

Example: Diet Problem

For each nutrient N , use a multiple C_N of the equation for that nutrient.

Can then add multiples of the constraint that you get a non-negative amount of each food.

Example: Diet Problem

Think of C_N as a cost of nutrient N . We pick values so that for each food item, f , we have

$$\text{Cost}(f) \geq \sum_N C_N \cdot (\text{Amount of nutrient } N \text{ in } f).$$

Example: Diet Problem

Think of C_N as a cost of nutrient N . We pick values so that for each food item, f , we have

$$\text{Cost}(f) \geq \sum_N C_N \cdot (\text{Amount of nutrient } N \text{ in } f).$$

Costs C_N to get a unit of nutrient N . This means total cost of a balanced diet is at least

$$\sum_N C_N \cdot (\text{Required amount of nutrient } N).$$

Observation

Note that if you want to actually obtain this lower bound, you cannot buy overpriced foods. Can only afford to buy foods with

$$\text{Cost}(f) = \sum_N C_N \cdot (\text{Amount of nutrient } N \text{ in } f).$$

Observation

Note that if you want to actually obtain this lower bound, you cannot buy overpriced foods. Can only afford to buy foods with

$$\text{Cost}(f) = \sum_N C_N \cdot (\text{Amount of nutrient } N \text{ in } f).$$

This is an example of a general phenomena called **complementary slackness**.

Complementary Slackness

Theorem

Consider a primal LP:

Minimize $v \cdot x$ subject to $Ax \geq b$,


and its dual LP:

Maximize $y \cdot b$ subject to $y^T A = v$, $y \geq 0$.

Then in the solutions, $y_i > 0$ only if the i^{th} equation in x is tight.


Problem

Assuming that the highlighted point is the optimum to the linear program below, which equations might have non-zero coefficients in the solution to the dual program?



Solution

Only 2 and 4.



Summary

- Every LP has dual LP.
- Solutions to dual bound solutions to primal.
- LP and dual have same answer!
- Complementary slackness.