

# Divide-and-Conquer: Master Theorem

Neil Rhodes

Department of Computer Science and Engineering  
University of California, San Diego

Algorithmic Design and Techniques  
Algorithms and Data Structures at edX

# Outline

- ① What is the Master Theorem
- ② Proof of Master Theorem

$$T(n) = T\left(\tfrac{n}{2}\right) + O(1)$$

$$T(n) = T\left(\frac{n}{2}\right) + O(1)$$



$$T(n) = O(\log n)$$

$$T(n) = 4\,T\left(\frac{n}{2}\right) + O(n)$$

$$T(n) = 4T\left(\frac{n}{2}\right) + O(n)$$



$$T(n) = O(n^2)$$

$$T(n) = 3T\left(\frac{n}{2}\right) + O(n)$$

$$T(n) = 3T\left(\frac{n}{2}\right) + O(n)$$



$$T(n) = O(n^{\log_2 3})$$

$$T(n) = 2T\left(\frac{n}{2}\right) + O(n)$$

$$T(n) = 2T\left(\frac{n}{2}\right) + O(n)$$



$$T(n) = O(n \log n)$$

# Master Theorem

## Theorem

If  $T(n) = aT\left(\lceil \frac{n}{b} \rceil\right) + O(n^d)$

# Master Theorem

## Theorem

If  $T(n) = aT\left(\lceil \frac{n}{b} \rceil\right) + O(n^d)$  (for constants  $a > 0, b > 1, d \geq 0$ ), then:

# Master Theorem

## Theorem

If  $T(n) = aT\left(\lceil \frac{n}{b} \rceil\right) + O(n^d)$  (for constants  $a > 0, b > 1, d \geq 0$ ), then:

$$T(n) = \begin{cases} \dots & \text{if } \dots \\ \dots & \text{if } \dots \\ \dots & \text{if } \dots \end{cases}$$

# Master Theorem

## Theorem

If  $T(n) = aT\left(\lceil \frac{n}{b} \rceil\right) + O(n^d)$  (for constants  $a > 0, b > 1, d \geq 0$ ), then:

$$T(n) = \begin{cases} O(n^d) & \text{if } d > \log_b a \\ & \dots \\ & \dots \end{cases}$$

# Master Theorem

## Theorem

If  $T(n) = aT\left(\lceil \frac{n}{b} \rceil\right) + O(n^d)$  (for constants  $a > 0, b > 1, d \geq 0$ ), then:

$$T(n) = \begin{cases} O(n^d) & \text{if } d > \log_b a \\ O(n^d \log n) & \text{if } d = \log_b a \end{cases}$$

# Master Theorem

## Theorem

If  $T(n) = aT\left(\lceil \frac{n}{b} \rceil\right) + O(n^d)$  (for constants  $a > 0, b > 1, d \geq 0$ ), then:

$$T(n) = \begin{cases} O(n^d) & \text{if } d > \log_b a \\ O(n^d \log n) & \text{if } d = \log_b a \\ O(n^{\log_b a}) & \text{if } d < \log_b a \end{cases}$$

# Master Theorem Example 1

$$T(n) = 4T\left(\frac{n}{2}\right) + O(n)$$

# Master Theorem Example 1

$$T(n) = 4 T\left(\frac{n}{2}\right) + O(n)$$

$$a = 4$$

# Master Theorem Example 1

$$T(n) = 4T\left(\frac{n}{2}\right) + O(n)$$

$$a = 4$$

$$b = 2$$

# Master Theorem Example 1

$$T(n) = 4T\left(\frac{n}{2}\right) + O(n^{\textcolor{red}{1}})$$

$$a = 4$$

$$b = 2$$

$$d = \textcolor{red}{1}$$

## Master Theorem Example 1

$$T(n) = 4T\left(\frac{n}{2}\right) + O(n)$$

$$a = 4$$

$$b = 2$$

$$d = 1$$

Since  $d < \log_b a$ ,  $T(n) = O(n^{\log_b a}) = O(n^2)$

## Master Theorem Example 2

$$T(n) = 3T\left(\frac{n}{2}\right) + O(n)$$

## Master Theorem Example 2

$$T(n) = 3T\left(\frac{n}{2}\right) + O(n)$$

$$a = 3$$

## Master Theorem Example 2

$$T(n) = 3T\left(\frac{n}{2}\right) + O(n)$$

$$a = 3$$

$$b = 2$$

## Master Theorem Example 2

$$T(n) = 3T\left(\frac{n}{2}\right) + O(n^{\textcolor{red}{1}})$$

$$a = 3$$

$$b = 2$$

$$\textcolor{red}{d} = 1$$

## Master Theorem Example 2

$$T(n) = 3T\left(\frac{n}{2}\right) + O(n)$$

$$a = 3$$

$$b = 2$$

$$d = 1$$

Since  $d < \log_b a$ ,

$$T(n) = O(n^{\log_b a}) = O(n^{\log_2 3})$$

## Master Theorem Example 3

$$T(n) = 2T\left(\frac{n}{2}\right) + O(n)$$

## Master Theorem Example 3

$$T(n) = 2T\left(\frac{n}{2}\right) + O(n)$$

$$a = 2$$

## Master Theorem Example 3

$$T(n) = 2T\left(\frac{n}{2}\right) + O(n)$$

$$a = 2$$

$$b = 2$$

## Master Theorem Example 3

$$T(n) = 2T\left(\frac{n}{2}\right) + O(n^{\textcolor{red}{1}})$$

$$a = 2$$

$$b = 2$$

$$d = \textcolor{red}{1}$$

## Master Theorem Example 3

$$T(n) = 2T\left(\frac{n}{2}\right) + O(n)$$

$$a = 2$$

$$b = 2$$

$$d = 1$$

Since  $d = \log_b a$ ,

$$T(n) = O(n^d \log n) = O(n \log n)$$

## Master Theorem Example 4

$$T(n) = T\left(\frac{n}{2}\right) + O(1)$$

## Master Theorem Example 4

$$T(n) = \textcolor{red}{1} T\left(\frac{n}{2}\right) + O(1)$$

$$a = \textcolor{red}{1}$$

## Master Theorem Example 4

$$T(n) = T\left(\frac{n}{2}\right) + O(1)$$

$$a = 1$$

$$b = 2$$

## Master Theorem Example 4

$$T(n) = T\left(\frac{n}{2}\right) + O(n^0)$$

$$a = 1$$

$$b = 2$$

$$d = 0$$

## Master Theorem Example 4

$$T(n) = T\left(\frac{n}{2}\right) + O(1)$$

$$a = 1$$

$$b = 2$$

$$d = 0$$

Since  $d = \log_b a$ ,  $T(n) = O(n^d \log n) = O(n^0 \log n) = O(\log n)$

## Master Theorem Example 5

$$T(n) = 2T\left(\frac{n}{2}\right) + O(n^2)$$

## Master Theorem Example 5

$$T(n) = 2T\left(\frac{n}{2}\right) + O(n^2)$$

$$a = 2$$

## Master Theorem Example 5

$$T(n) = 2T\left(\frac{n}{2}\right) + O(n^2)$$

$$a = 2$$

$$b = 2$$

## Master Theorem Example 5

$$T(n) = 2T\left(\frac{n}{2}\right) + O(n^2)$$

$$a = 2$$

$$b = 2$$

$$d = 2$$

## Master Theorem Example 5

$$T(n) = 2T\left(\frac{n}{2}\right) + O(n^2)$$

$$a = 2$$

$$b = 2$$

$$d = 2$$

Since  $d > \log_b a$ ,  $T(n) = O(n^d) = O(n^2)$

# Outline

- 1 What is the Master Theorem
- 2 Proof of Master Theorem

# Master Theorem

## Theorem

If  $T(n) = aT\left(\lceil \frac{n}{b} \rceil\right) + O(n^d)$  (for constants  $a > 0, b > 1, d \geq 0$ ), then:


$$T(n) = \begin{cases} O(n^d) & \text{if } d > \log_b a \\ O(n^d \log n) & \text{if } d = \log_b a \\ O(n^{\log_b a}) & \text{if } d < \log_b a \end{cases}$$

$$T(n) = a T\left(\left\lceil \frac{n}{b} \right\rceil\right) + O(n^d)$$

$$\boxed{n}$$

$$T(n) = aT\left(\lceil \frac{n}{b} \rceil\right) + O(n^d)$$

level




$$T(n) = aT\left(\left\lceil \frac{n}{b} \right\rceil\right) + O(n^d)$$

level

0

1



$$T(n) = aT\left(\left\lceil \frac{n}{b} \right\rceil\right) + O(n^d)$$


level

0


1

⋮


$i$




$$T(n) = aT\left(\left\lceil \frac{n}{b} \right\rceil\right) + O(n^d)$$




$$T(n) = aT\left(\left\lceil \frac{n}{b} \right\rceil\right) + O(n^d)$$




$$T(n) = aT\left(\left\lceil \frac{n}{b} \right\rceil\right) + O(n^d)$$




$$T(n) = aT\left(\left\lceil \frac{n}{b} \right\rceil\right) + O(n^d)$$




$$T(n) = aT\left(\lceil \frac{n}{b} \rceil\right) + O(n^d)$$




$$T(n) = aT\left(\left\lceil \frac{n}{b} \right\rceil\right) + O(n^d)$$



$$T(n) = aT\left(\left\lceil \frac{n}{b} \right\rceil\right) + O(n^d)$$




$$T(n) = aT\left(\lceil \frac{n}{b} \rceil\right) + O(n^d)$$



$$T(n) = aT\left(\lceil \frac{n}{b} \rceil\right) + O(n^d)$$

level

0



work

$$O(n^d)$$

$$aO\left(\frac{n}{b}\right)^d = O(n^d)\frac{a}{b^d}$$

⋮

$i$

$$\cdots \boxed{n/b^i} \cdots$$

$$a^i O\left(\frac{n}{b^i}\right)^d = O(n^d)\left(\frac{a}{b^d}\right)^i$$

⋮

log<sub>b</sub> n

$$\boxed{1}$$

⋮

$$\boxed{1}$$


⋮

$$a^{\log_b n}$$

$$T(n) = aT\left(\lceil \frac{n}{b} \rceil\right) + O(n^d)$$

level

0



1

⋮

$i$

$\cdots [n/b^i] \cdots$

⋮

$\log_b n$



work

$$O(n^d)$$

$$aO\left(\frac{n}{b}\right)^d = O(n^d)\frac{a}{b^d}$$

⋮

$$a^i O\left(\frac{n}{b^i}\right)^d = O(n^d)\left(\frac{a}{b^d}\right)^i$$


⋮

$$a^{\log_b n} = O(n^{\log_b a})$$

$$T(n) = aT\left(\left\lceil \frac{n}{b} \right\rceil\right) + O(n^d)$$

level

0



1

⋮

$i$

⋮

$\log_b n$

work

$$O(n^d)$$

$$aO\left(\frac{n}{b}\right)^d = O(n^d)\left(\frac{a}{b^d}\right)$$

⋮

$$a^i O\left(\frac{n}{b^i}\right)^d = O(n^d)\left(\frac{a}{b^d}\right)^i$$

⋮

$$a^{\log_b n} = O(n^{\log_b a})$$

$$\text{Total: } \sum_{i=0}^{\log_b n} O(n^d)\left(\frac{a}{b^d}\right)^i$$

# Geometric Series

For  $r \neq 1$ :

$$a + ar + ar^2 + ar^3 + \cdots + ar^{n-1}$$

# Geometric Series

For  $r \neq 1$ :

$$\begin{aligned} & a + ar + ar^2 + ar^3 + \cdots + ar^{n-1} \\ &= a \frac{1 - r^n}{1 - r} \end{aligned}$$

# Geometric Series

For  $r \neq 1$ :

$$\begin{aligned} & a + ar + ar^2 + ar^3 + \cdots + ar^{n-1} \\ &= a \frac{1 - r^n}{1 - r} \\ &= \left\{ \right. \end{aligned}$$

# Geometric Series

For  $r \neq 1$ :

$$\begin{aligned} & a + ar + ar^2 + ar^3 + \cdots + ar^{n-1} \\ &= a \frac{1 - r^n}{1 - r} \\ &= \begin{cases} O(a) & \text{if } r < 1 \end{cases} \end{aligned}$$

# Geometric Series

For  $r \neq 1$ :

$$\begin{aligned} & a + ar + ar^2 + ar^3 + \cdots + ar^{n-1} \\ &= a \frac{1 - r^n}{1 - r} \\ &= \begin{cases} O(a) & \text{if } r < 1 \\ O(ar^{n-1}) & \text{if } r > 1 \end{cases} \end{aligned}$$

Case 1:  $\frac{a}{b^d} < 1$  ( $d > \log_b a$ )

$$\sum_{i=0}^{\log_b n} O(n^d) \left(\frac{a}{b^d}\right)^i$$

Case 1:  $\frac{a}{b^d} < 1$  ( $d > \log_b a$ )

$$\begin{aligned}& \sum_{i=0}^{\log_b n} O(n^d) \left(\frac{a}{b^d}\right)^i \\&= O(n^d)\end{aligned}$$

Case 2:  $\frac{a}{b^d} = 1$  ( $d = \log_b a$ )

$$\sum_{i=0}^{\log_b n} O(n^d) \left(\frac{a}{b^d}\right)^i$$

Case 2:  $\frac{a}{b^d} = 1$  ( $d = \log_b a$ )

$$\begin{aligned}& \sum_{i=0}^{\log_b n} O(n^d) \left(\frac{a}{b^d}\right)^i \\&= \sum_{i=0}^{\log_b n} O(n^d)\end{aligned}$$

Case 2:  $\frac{a}{b^d} = 1$  ( $d = \log_b a$ )

$$\begin{aligned}& \sum_{i=0}^{\log_b n} O(n^d) \left(\frac{a}{b^d}\right)^i \\&= \sum_{i=0}^{\log_b n} O(n^d) \\&= (1 + \log_b n) O(n^d)\end{aligned}$$

Case 2:  $\frac{a}{b^d} = 1$  ( $d = \log_b a$ )

$$\begin{aligned}& \sum_{i=0}^{\log_b n} O(n^d) \left(\frac{a}{b^d}\right)^i \\&= \sum_{i=0}^{\log_b n} O(n^d) \\&= (1 + \log_b n) O(n^d) \\&= O(n^d \log n)\end{aligned}$$

Case 3:  $\frac{a}{b^d} > 1$  ( $d < \log_b a$ )

$$\sum_{i=0}^{\log_b n} O(n^d) \left(\frac{a}{b^d}\right)^i$$

Case 3:  $\frac{a}{b^d} > 1$  ( $d < \log_b a$ )

$$\begin{aligned}& \sum_{i=0}^{\log_b n} O(n^d) \left(\frac{a}{b^d}\right)^i \\&= O\left(O(n^d) \left(\frac{a}{b^d}\right)^{\log_b n}\right)\end{aligned}$$

Case 3:  $\frac{a}{b^d} > 1$  ( $d < \log_b a$ )

$$\begin{aligned}& \sum_{i=0}^{\log_b n} O(n^d) \left(\frac{a}{b^d}\right)^i \\&= O\left(O(n^d) \left(\frac{a}{b^d}\right)^{\log_b n}\right) \\&= O\left(O(n^d) \frac{a^{\log_b n}}{b^{d \log_b n}}\right)\end{aligned}$$

Case 3:  $\frac{a}{b^d} > 1$  ( $d < \log_b a$ )

$$\begin{aligned}& \sum_{i=0}^{\log_b n} O(n^d) \left(\frac{a}{b^d}\right)^i \\&= O\left(O(n^d) \left(\frac{a}{b^d}\right)^{\log_b n}\right) \\&= O\left(O(n^d) \frac{a^{\log_b n}}{b^{d \log_b n}}\right) \\&= O\left(O(n^d) \frac{n^{\log_b a}}{n^d}\right)\end{aligned}$$

Case 3:  $\frac{a}{b^d} > 1$  ( $d < \log_b a$ )

$$\begin{aligned}& \sum_{i=0}^{\log_b n} O(n^d) \left(\frac{a}{b^d}\right)^i \\&= O\left(O(n^d) \left(\frac{a}{b^d}\right)^{\log_b n}\right) \\&= O\left(O(n^d) \frac{a^{\log_b n}}{b^{d \log_b n}}\right) \\&= O\left(O(n^d) \frac{n^{\log_b a}}{n^d}\right) \\&= O(n^{\log_b a})\end{aligned}$$

# Summary

Master theorem is a shortcut:

# Summary

Master theorem is a shortcut:

## Theorem

If  $T(n) = aT\left(\lceil \frac{n}{b} \rceil\right) + O(n^d)$  (for constants  $a > 0, b > 1, d \geq 0$ ), then:

$$T(n) = \begin{cases} O(n^d) & \text{if } d > \log_b a \\ O(n^d \log n) & \text{if } d = \log_b a \\ O(n^{\log_b a}) & \text{if } d < \log_b a \end{cases}$$